

Fog and Thunderstorm Forecasting in Melbourne, Australia

Harvey Stern*

Bureau of Meteorology, Australia

1. Introduction

A twelve-month "real-time" trial of a methodology utilised to generate Day-1 to Day-7 forecasts, by mechanically integrating judgmental (human) and automated predictions, was conducted between 20 August 2005 and 19 August 2006. After 365 days, the trial revealed that, overall, the various components (rainfall amount, sensible weather, minimum temperature, and maximum temperature) of Melbourne forecasts so generated explained 41.3% variance of the weather, 7.9% more variance than the 33.4% variance explained by the human (official) forecasts alone (Stern, 2007a, 2007b).

The trial continues and the purpose of the current paper is to report on its performance at predicting fog and thunderstorms between August 2005 and June 2007, a period of almost two years.

2. Verification of fog forecasts

With regard to the accuracy of forecasts of fog, for verification purposes, it is said that there has been fog in the metropolitan area during a particular day when at least one of the 0300, 0600, 0900, 1200, 1500, 1800, 2100, or 2400 Melbourne CBD and/or Melbourne Airport observations include a report of fog (including shallow fog) and/or distant fog.

The automated component of the system used to forecast fog is that described by Stern and Parkyn, 1998, 1999, 2000, 2001). This component is a logistic model that, in summary, feeds observational data from the preceding afternoon into a set of prediction equations, developed by applying logistic regression to sets of synoptically stratified data, to yield an estimate of the probability of fog.

The combining process was shown to lift the Critical Success Index (CSI) (Wilks, 1995) from 13.6% to 15.2%, and to lift the Probability of Detection (PoD) from 17.7% to 28.0. However, the lift in the CSI and PoD for fog forecasts was achieved at a cost of a corresponding increase in the False Alarm Ratios (FARs), from 63.2% to 74.9%.

Although a lift did not occur in every single instance when the verification data was analysed with all lead times taken separately, a lift occurred in most instances. The exceptions were in the cases of Day-1 and Day-2 forecasts of fog, where the CSIs were substantially below corresponding CSIs for the human (official) forecasts.

These forecasts are worthy of comment. The inability (of the combining process) to improve on the Day-1 and Day-2 official forecasts of fog may very well be a consequence of the effort that the forecasting personnel of the Victorian Regional Office (and others) have invested over the years into short term fog and low cloud forecasting at Melbourne Airport (Goodhead, 1978; Keith, 1978; Stern and Parkyn, 1998, 1999, 2000, 2001; Newham, 2004, Weymouth et al, 2007; Newham et al, 2007), most recently using a Bayesian network to combine various components of forecasting guidance.

This effort may have resulted in such a high level of pre-existing human forecast skill at short-term predicting of fog, that mechanically combining human fog forecasts with automated fog forecasts (generated by a methodology more than five years old) actually caused a decline in accuracy.

3. Verification of thunderstorm forecasts

With regard to the accuracy of forecasts of thunderstorms, for verification purposes, it is said that there has been a thunderstorm in the metropolitan area during a particular day when at least one of the 0300, 0600, 0900, 1200, 1500, 1800, 2100, or 2400 Melbourne CBD and/or Melbourne Airport observations include a report of cumulonimbus with an anvil and/or lightning and/or funnel cloud and/or thunder (with or without precipitation).

The automated component of the system used to forecast thunderstorms is that described by Stern (2004). This component is a logistic model that, in summary, feeds a Quantitative Precipitation Forecast (QPF), and a Probability of Precipitation (PoP) estimate, into a set of prediction equations, developed by applying logistic regression to sets of synoptically stratified data, to yield an estimate of the probability of thunderstorms.

The combining process was shown to lift the Critical Success Index (CSI) (Wilks, 1995) from 13.7% to 18.2%, and to lift the Probability of Detection (PoD) from 15.9% to 26.7%. However, the lift in the CSI and PoD for thunderstorm forecasts was achieved at a cost of a corresponding increase in the False Alarm Ratios (FARs), from 50.9% to 63.5%. A lift in CSI occurred in every instance when the verification data

was analysed with lead times taken separately, except for Day-2

4. Concluding Remark

The data presented demonstrate that, overall, adopting a strategy of combining human (official) and automated predictions of fog and thunderstorms enhances the skill displayed by such predictions.

However, the data suggest that predictions, which have been prepared by operational meteorologists armed with the very latest techniques, are sometimes capable of outperforming forecasts generated by combining those predictions with automated predictions that have been produced utilising old techniques.

5. References

- Goodhead, H. M., 1978: An objective method for predicting fog and mist at Melbourne Airport. *Airmet Conference*, 9-10 Feb., 1978, Canberra Meteor. Soc., Roy. Meteor. Soc. (Aust. Branch), Canberra, Australia.
- Keith, R., 1978: Formation of low cloud at Melbourne Airport in a lee trough. *Airmet Conference*, 9-10 Feb., 1978, Canberra Meteor. Soc., Roy. Meteor. Soc. (Aust. Branch), Canberra, Australia, 39-44.
- Newham, P., 2004: Fog forecasting for Melbourne Airport. *3rd Conference on Fog, Fog Collection, and Dew*, Cape Town, South Africa, 11-15 October, 2001.
- Newham, P., Boneh, T., Weymouth, G. T., Potts, R., Bally, J., Nicholson, A., and Korb, K. Dealing with uncertainty in fog forecasting for major airports in Australia. *4th Conference on Fog, Fog Collection and Dew*, La Serena, Chile, 22-27 Jul., 2007.
- Stern, H., 2004: Using a knowledge based system to predict thunderstorms. Presented at *International Conference on Storms, Storms Science to Disaster Mitigation*, Brisbane, Queensland, Australia 5-9 Jul., 2004.
- Stern, H., 2007a: Increasing weather forecast accuracy by mechanically combining human and automated predictions using a knowledge based system. *23rd Conference on Interactive Information and Processing Systems, San Antonio, Texas, USA 14-18 Jan., 2007*.
- Stern, H., 2007b: Improving forecasts with mechanically combined predictions. *Bulletin of the American Meteorological Society (BAMS)*, June 2007, 88:850-851.
- Stern, H. and Parkyn, K., 1998: Synoptic climatology of fog at Melbourne Airport. Abstracts, *ANZ Climate Forum*, Perth, Australia, 1998, 51.
- Stern, H. and Parkyn, K., 1999: Predicting the likelihood of fog at Melbourne Airport, *8th Conference on Aviation, Range and Aerospace Meteorology*, Amer. Meteor. Soc., Dallas, Texas, 10-15 Jan., 1999.
- Stern, H. and Parkyn, K., 2000: Low cloud at Melbourne Airport: A synoptic climatology leading to a forecasting technique, *AMOS 2000, incorporating 7th National AMOS conference & 5th Australasian conference on the Physics of Remote Sensing of Atmosphere and Ocean*, The University of Melbourne, Melbourne, 7-9 February 2000.
- Stern, H. and Parkyn, K. 2001: A web-based Melbourne Airport fog and low cloud forecasting technique. *2nd Conference on Fog and Fog Collection*, St John's, New Foundland, Canada 15-20 Jul., 2001.
- Weymouth, G. T., Boneh, T., Newham, P., Bally, J., Potts, R., and Korb, K. Dealing with uncertainty in fog forecasting for major airports in Australia. *4th Conference on Fog, Fog Collection and Dew*, La Serena, Chile, 22-27 Jul., 2007.
- Wilks, D., 1995: Statistical methods in atmospheric sciences. *Academic Press*.




Fig 1 CSIs for fog forecasts




Fig 2 PoDs for fog forecasts




Fig 3 FARs for fog forecasts




Fig 4 CSIs for thunderstorm forecasts




Fig 5 PoDs for thunderstorm forecasts




Fig 6 FARs for thunderstorm forecasts